

A NEW QUANTUM PHASE IN TWO DIMENSIONS

GIULIANO BENENTI, XAVIER WAIN TAL, AND JEAN-LOUIS PICHARD

*CEA, Service de Physique de l'Etat Condensé,
Centre d'Etudes de Saclay, F-91191 Gif-sur-Yvette, France*

For intermediate Coulomb energy to Fermi energy ratios r_s , spinless fermions in a random potential form a new quantum phase which is different from the Fermi glass and the Wigner crystal. From a numerical study of small clusters, we show that this phase is characterized by an ordered flow of enhanced persistent currents and occurs for r_s values where a metallic phase has been observed in two dimensions.

1 Introduction

A crucial parameter for a system of charged particles is the Coulomb energy to Fermi energy ratio r_s . In a disordered two-dimensional system, the ground state is obvious in the limiting cases. For large r_s , the charges form a pinned Wigner crystal, the Coulomb repulsion being dominant over the kinetic energy and the disorder. For small r_s , the interaction becomes negligible and the ground state is a Fermi glass with Anderson localized one electron states. For intermediate r_s , a lot of transport measurements following the pioneering works of Kravchenko and co-workers¹ and made with electron and hole gases give evidence of a metallic phase in two dimensions, observed² for instance when $6 < r_s < 9$ for a hole gas in GaAs heterostructures. In a recent paper³ we have shown that spinless fermions with Coulomb repulsion in disordered $2d$ clusters exhibits, for a similar range of r_s values, a new ground state characterized by an ordered flow of enhanced persistent currents. A study over the statistical ensemble of the currents supported by the ground state gives us two well defined values r_s^F and r_s^W bounding this new phase.

2 The model

We consider a simple model of $N = 4$ Coulomb interacting spinless fermions in a random potential defined on a square lattice with $L^2 = 36$ sites. The Hamiltonian reads:

$$H = -t \sum_{\langle i,j \rangle} c_i^\dagger c_j + \sum_i v_i n_i + U \sum_{i \neq j} \frac{n_i n_j}{2r_{ij}}, \quad (1)$$

where $c_i^\dagger$ (c_i) creates (destroys) an electron in the site i , t is the strength of the hopping terms between nearest neighbours ($t = 1$ in the following) and r_{ij} is the inter-particle distance for a $2d$ torus. The random potential v_i of the site i with occupation number $n_i = c_i^\dagger c_i$ is taken from a box distribution of width W . The interaction strength U yields a Coulomb energy to Fermi energy ratio $r_s = U/(2t\sqrt{\pi n_e})$ for a filling factor $n_e = N/L^2$. A Fermi golden rule approximation for the elastic scattering time leads to $k_F l \approx 192\pi n_e (t/W)^2$. Here k_F and l denote the Fermi wave vector and the elastic mean free path respectively and the above estimate is valid for filling factors $n_e \ll 1$. We consider disorder strengths $W = 5, 10, 15$ corresponding to $k_F l = 2.7, 0.7$ and 0.3 respectively. The boundary conditions are always taken periodic in the transverse y -direction, and such that the system encloses an Aharonov-Bohm flux ϕ in the longitudinal x -direction. Imposing $\phi = \pi/2$ ($\phi = \pi$ corresponds to anti-periodic boundary conditions), one drives a persistent current of total longitudinal and transverse components given by

$$I_l = -\frac{\partial E}{\partial \phi} = \frac{\sum_i I_i^l}{L} \quad \text{and} \quad I_t = \frac{\sum_i I_i^t}{L}. \quad (2)$$




Figure 1: Behavior of a single cluster for $W = 15$ (left column) and $W = 5$ (right column) as a function of r_s . From top to bottom: low energy spectrum (a $1.9r_s$ term has been subtracted); first energy spacing Δ_0 ; crystallization parameter γ for the ground state (filled diamonds) and for the first excited state (open diamonds); longitudinal I_l (empty circles) and transverse I_t (empty triangles) current (left scale) and number of occupied sites ξ_s (filled circles, right scale).




Figure 2: Local currents for the sample of Fig. 1 at $W = 5$, $r_s = 0$ (left) and $r_s = 17$ (right).

The local current I_i^l flowing at the site i in the longitudinal direction is defined by $I_i^l = 2\text{Im}(\langle \Psi_0 | c_{i_{x+1}, i_y}^\dagger c_{i_x, i_y} | \Psi_0 \rangle)$, with an analogous expression for I_i^t . The response is paramagnetic if $I_l > 0$ and diamagnetic if $I_l < 0$.

3 Numerical results

Fig. 1 exhibits behaviours characteristic of individual small clusters, comparing the strongly disordered regime $k_F l < 1$ ($W = 15$) with the regime $k_F l > 1$ ($W = 5$). For $k_F l < 1$, looking at the low energy part of the spectrum one can see that, as we gradually turn on the interaction, classification of the levels remains invariant up to first avoided crossings, where a Landau theory of the Fermi glass is certainly no longer possible. The electronic density $\rho_i = \langle \Psi_0 | n_i | \Psi_0 \rangle$ of the ground state $|\Psi_0\rangle$ is maximum in the minima of the site potentials for the Fermi glass. After the second avoided crossing, ρ_i is negligible except for four sites forming a lattice of charges as close as possible to the Wigner crystal triangular network in the imposed square lattice. The degeneracy of the crystal is removed by the disorder, the array being pinned in 4 sites of favourable energies. For the same cluster, we have calculated $C(r) = N^{-1} \sum_i \rho_i \rho_{i+r}$ and used the parameter $\gamma = \max_r C(r) - \min_r C(r)$ to characterize the melting of the crystal ($\gamma = 1$ for a crystal and 0 for a liquid). Calculated for the ground state and the first excited state, γ allows us to identify the second crossing with the melting of the crystal. Moreover, one can see that the crystal becomes unstable in the intermediate phase, while the ground state is related to the first excitation of the crystal. Around the crossings, the longitudinal current I_l and the participation ratio $\xi_s = N^2(\sum_i \rho_i^2)^{-1}$ of the ground state (i.e. the number of sites that it occupies) are enhanced. The general picture reminds us of strongly disordered chains⁴ where level crossings are associated to a charge reorganization of the ground state and comes with an enhancement of the persistent current. For $k_F l \geq 1$ ($W = 5$), the previous level crossings are now almost suppressed by a stronger level repulsion and charge crystallization occurs more continuously. There is instead a broad enhancement of I_l which, in contrast to the regime $k_F l < 1$, is not accompanied by a corresponding increase in the number of occupied sites ξ_s . One can also notice that, increasing r_s , I_t is suppressed before I_l . This can be understood looking at the local currents (see Fig. 2): the interaction causes the transition from a disordered array of currents making closed loops to a plastic flow of correlated currents in the longitudinal direction⁵.

We summarize in Fig. 3 results from a statistical study of an ensemble of 10^3 clusters for $W = 5, 10, 15$. One can see an interaction-induced enhancement of the averaged persistent




Figure 3: Statistical study of an ensemble of 10^3 clusters for $W = 5$ (circles), 10 (squares) and 15 (triangles) as a function of r_s . Mean value of the longitudinal current $\langle I_l \rangle$. Fraction C_d of diamagnetic samples. Distribution of the logarithms of the paramagnetic currents $P(\text{Ln}|I_{l,p}|)$ and of the diamagnetic currents $P(\text{Ln}|I_{l,d}|)$ at $r_s = 1.7$ and $W = 15$ (for an ensemble of 10^4 clusters), fitted by log-normal functions of mean $\mu = -5.8$ and variance $\sigma^2 = 2.3$ in the paramagnetic case, $\mu = -6.5$ and $\sigma^2 = 2.9$ in the diamagnetic case. Log-averages of the longitudinal paramagnetic (empty symbols) and diamagnetic (filled symbols) currents. The straight lines are exponential fits giving $r_s^W = 9.5$ ($W = 5$), $r_s^W = 10.3$ ($W = 10$), $r_s^W = 12.4$ ($W = 15$). Log averages of the transverse currents; the exponential fits give $r_s^F = 3.3$ ($W = 5$), $r_s^F = 3.7$ ($W = 10$), $r_s^F = 4.5$ ($W = 15$). Mean number of occupied sites $\langle \xi_s \rangle$. Mean crystallization parameter $\langle \gamma \rangle$.




Figure 4: Sketch of the phase diagram for 2d spinless fermions in a random potential, with r_s^W (filled circle) and r_s^F (empty circle) values obtained from Fig. 3.

current $\langle I_l \rangle$ by almost one order of magnitude when $r_s \approx 7$ and $W = 5$. This effect can be partially explained by the suppression of the fraction of diamagnetic clusters c_d . The paramagnetic $I_{l,p}$ and diamagnetic $I_{l,d}$ longitudinal currents, and $|I_t|$ have acceptable log-normal distributions for all values of r_s when $W \geq 5$. The stronger is the disorder, the better is the log-normal shape of the distributions (see Fig. 3 for $W = 15$). The averages of the logarithms of $I_{l,p}$, $-I_{l,d}$ and $|I_t|$ give the typical values shown in Fig. 3. They exponentially decay as $|I_t| \propto \exp(-r_s/r_s^F)$ and $I_l \propto \exp(-r_s/r_s^W)$ when r_s is large enough. The values of r_s^F and r_s^W extracted from the exponential fits (straight lines of Fig. 3) are given in Fig. 4, where a sketch of the phase diagram is proposed.

4 Conclusions

A simple model of Coulomb interacting spinless fermions in a random potential allows us to identify a new quantum phase, clearly separated from the Fermi glass and the Wigner crystal. This phase is characterized by a plastic flow of currents and a response to an Aharonov-Bohm flux with a sign independent of the microscopic disorder. For $k_F l \geq 1$, I_l displays a strong enhancement for intermediate r_s values, which could be the signature of a new metal in the thermodynamic limit. For $k_F l < 1$, ξ_s and I_l convey similar information while the increase of I_l is accompanied by a decrease of ξ_s and an increase of γ when $k_F l > 1$ (see Fig. 3). This suggests that, in the latter case, transport in the new phase results from a collective motion of charges and not from a delocalization of individual charges.

This work is partially supported by a TMR network of the EU.

References

1. S.V. Kravchenko *et al.* *Phys. Rev. B* **50**, 8039 (1994); **51**, 7038 (1995).
2. A.R. Hamilton, M.Y. Simmons, M. Pepper, E.H. Linfield, P.D. Rose and D.A. Ritchie, *Phys. Rev. Lett.* **82**, 1542 (1999).
3. G. Benenti, X. Waintal and J.-L. Pichard, cond-mat/9904096.
4. P. Schmitteckert, R.A. Jalabert, D. Weinmann and J.-L. Pichard, *Phys. Rev. Lett.* **81**, 2308 (1998) (see also in this volume).
5. R. Berkovits and Y. Avishai, *Phys. Rev. B* **57**, R15076 (1998).